

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 338 209 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

27.08.2003 Bulletin 2003/35

(51) Int Cl.7: **A23L 1/16**

(21) Application number: **02425101.9**

(22) Date of filing: **26.02.2002**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

Designated Extension States:

AL LT LV MK RO SI

• **Bulat, Irene**

43100 Parma (IT)

• **Caselli, Oreste**

43100 Parma (IT)

• **Mangiavacca, Nicola**

43100 Parma (IT)

(71) Applicant: **BARILLA ALIMENTARE S.P.A.**

43100 Parma (IT)

(74) Representative: **Zambardino, Umberto et al**

Botti & Ferrari S.r.l.

Via Locatelli, 5

20124 Milano (IT)

(72) Inventors:

• **Dameno, Franco**

43100 Parma (IT)

(54) **A process for the preparation of cooked pasta, in particular for the preparation of ready-to-eat meals having a long shelf-life**

(57) A method is disclosed in which fresh alimentary pasta is prepared from a starting dough that has been acidified with an edible acid. The pasta is then dried to a moisture content of about 11%, cooked in salted water, strained, rinsed, dipped in an aqueous solution of an ed-

ible acid to achieve a pH no more than 4.1, and eventually metered into suitable packages and subjected to sanitization.

EP 1 338 209 A1

Description

Field of Application

[0001] The present invention broadly relates to the technological field of the production of ready-to-eat food dishes based on pre-cooked alimentary pasta.

[0002] In particular, the invention relates to a method of producing cooked (or pre-cooked) alimentary pasta, wherein the pasta is processed with an edible acid, and thus referred to as "acidic pasta", and is useful to prepare ready-to-eat dishes featuring long shelf-life at both room and refrigerator temperatures.

Prior Art

[0003] It is recognised that many types of cooked alimentary pasta that currently enter the commercial production of ready-to-eat dishes are the outcome of research efforts directed to make, on industrial basis, alimentary pasta for direct consumption which, on the one side, would have good organoleptic and quality characteristics, and on the other, could be preserved at room and refrigerator temperatures for long periods.

[0004] These efforts yielded for the most part techniques of producing alimentary cooked pasta, which include processing with an edible acid to reduce the pH of cooked pasta for substantially preventing the bacterial growth, since the high moisture content of cooked pasta provides good foundation for bacteria growth. It is on account of the added acid that cooked pasta so prepared has come to be commonly termed "acidic pasta".

[0005] One method of producing acidic pasta as above comprises adding an edible acid to the starting dough, shaping the newly made pasta (moisture content of 30 to 32%), and then cooking in water the pasta in the shape given to it. This method is described in US Patent 4,597,976, for example.

[0006] More particularly, US Patent 4,597,976 discloses a method of preparing ready-to-eat dishes of pre-cooked acidic pasta that also include flowing-state sauce. In this method, each component of the dish is processed with an edible acid so as to have a balanced pH of less than 4.6.

[0007] The pasta is obtained from a dough comprising flour, gluten, powder eggs, and an acid for alimentary use. The newly made pasta is then boiled in water, rinsed, strained, and steamed for 5 minutes at a temperature of at least 85°C. The cooked acidic pasta thus obtained is packaged in metered amounts, a sauce with a pH below 4.6 is added into the packages, and the packages are sealed and sterilized.

[0008] Such methods have two drawbacks:

- the presence of acid in the starting dough, as well as in the newly made pasta, causes a substantial destructure of the protein and starch matrix such that, after boiling in water, a cooked pasta hav-

ing low organoleptic properties is obtained, in particular a cooked pasta having low consistency or even an overcooked texture; and

- cooking in water causes acidic pasta to gradually lose the acid added to the starting dough, consequently cooked pasta has an acidity low enough to impair its preservability.

[0009] According to another known method of producing cooked pasta, no acid is added in the dough, and the pasta formed from the dough is cooked in water acidified with an acid.

[0010] This method has a serious drawback in that the distribution of the alimentary acid is not uniform throughout the structure of cooked pasta, but rather acidification is only adequate at the surface of cooked pasta.

[0011] Therefore, cooked acidic pasta prepared by the above method has a definitely shorter shelf-life.

[0012] The technical problem underlying the present invention is to provide a method of producing cooked acidic pasta, suitable for direct consumption or inclusion in ready-to-eat dishes, which has good organoleptic and textural characteristics and long shelf-life, thereby overcoming the aforementioned drawbacks of the prior art.

Summary of the Invention

[0013] The technical problem is solved by a method of producing cooked acidic alimentary pasta, characterized in that it comprises the steps of:

preparing an alimentary dough comprising semolina, water and at least one edible acid;

forming said alimentary dough obtaining fresh acidic pasta having a pH of 3.8 to 4.3 and a moisture content of 30 to 32%;

drying said fresh acidic pasta to a moisture content of 11.0 to 12.5%;

substantially completely cooking said dried pasta obtaining cooked acidic pasta;

treating said cooked acidic pasta with an aqueous solution of an alimentary acid to obtain cooked acidic pasta having a pH not more than 4.1; and

packaging said cooked acidic pasta and subjecting the packaged pasta to thermal sanitization.

[0014] The advantages of this invention accrue from that by applying the acid in two steps (to the starting dough and the cooked pasta) separated by a drying step, a cooked acidic pasta is obtained which achieves excellent organoleptic and textural characteristics, fully comparable to those of traditional freshly cooked pasta,

and this even at the end of long storage periods at room or refrigerator temperatures.

[0015] In particular, it is believed that acidification of the dough causes an optimal and uniform distribution of the edible acid down to the core of the fresh pasta prepared from it; that the intervening drying step, causing the gluten contained in the fresh pasta to reticulate, is effective to eliminate any destructuring effect of the acid on the structure of said fresh pasta; and that the second acidifying step carried out on the cooked pasta restores it to desired acidity values, which the cooking process is likely to have altered somewhat.

[0016] The starting dough may be any dough based on semolina and water, and comprising a standard edible acid for use in alimentary acidic pasta.

[0017] Preferably, gluten is added to the dough ingredients in amounts of 4 to 10% by weight on the dough weight, the preferred amount being 9%.

[0018] Gluten enhances the protein content of pasta and makes for a firmer and more consistent end product. Preferably, the moisture content of the dough is in the range of 30 to 40%.

[0019] The edible acid may be any organic or inorganic acid that is suitable for food applications and substantially soluble in water. Preferably, the edible acid is selected from a group comprising citric acid, fumaric acid, lactic acid, maleic acid, tartaric acid, sulphuric acid, hydrochloric acid, glucone-delta-lactone acid, and phosphoric acids. The preferred acid is glucone-delta-lactone acid.

[0020] The edible acid is added into the dough in varying amounts between 0.8 and 1.5% by weight on dough weight, 1% being the preferred amount.

[0021] The alimentary dough is formed conventionally.

[0022] The pasta is dried conventionally through conventional driers to yield dried pasta with a relative moisture content of up to 12.5%, preferably of 11 to 11.5%.

[0023] The alimentary pasta may be cooked in hot or boiling water in a substantial complete way for the time required to achieve desired characteristics, especially as regards firmness and consistency, depending on end users' preferences and demands.

[0024] Preferably, the pasta is cooked as required to achieve a so-called "al dente" cooking. Pasta is said to be cooked "al dente" when it is soft at the surface and firm inside. The moisture content of such pasta would usually lie between 53 and 63%, and the pasta itself shows firmness, consistency or texture, stickiness and accumulation/suppleness properties on the plate which are most appreciated by the consumer.

[0025] Preferably, the acidic pasta is rinsed with water on completion of the cooking step and before it goes through the second acidifying step.

[0026] Acidification is applied preferably by dipping the cooked acidic pasta into an aqueous solution of the alimentary acid of choice for a predetermined time to yield cooked acidic pasta having a pH no higher than

4.1, preferably in the range of 3.8 to 4.3.

[0027] Preferably, the pH of the aqueous solution should be 2.2 to 2.8.

[0028] Advantageously, the acidifying process of the cooked acidic pasta should be carried out at moderate temperatures of 18° to 25°C, preferably at room temperature, so as to retain the good organoleptic and textural characteristics of the cooked acidic pasta.

[0029] Following the acidifying process, the cooked acidic pasta is dried and loaded into packages, generally trays, in metered amounts. The package would be a gas-impermeable sealed type suitable for vacuum packaging or packaging under modified atmosphere.

[0030] Before packaging, the cooked acidic pasta may be additivated with an alimentary oil to attenuate the natural stickiness of pasta.

[0031] The modification of the atmosphere inside the package may be carried out by applying vacuum and optionally introducing into the package an inert gas such as nitrogen or carbon dioxide.

[0032] The packaged pasta must be subjected to sanitization for adequate preservation. This is done by subjecting the cooked pasta sealed into packages to a pasteurization and/or sterilization process known per se. For example, the packages containing cooked pasta can be pasteurized at a temperature of 80° to 90°C, by heating with steam or dipping in boiling water. A preferred method of pasteurizing or sterilizing cooked pasta includes exposure to microwave

[0033] The cooked acidic pasta obtained with the method of this invention has excellent organoleptic and textural characteristics that are fully comparable to those of traditional freshly cooked pasta. It can be consumed as such or can be further treated as desired before consumption. For example, before consumption cooked acidic pasta obtained with the method of this invention may be heated or fried with or without condiment added, while substantially maintaining its original characteristics.

[0034] In addition, the shelf-life of this pasta is made quite long because the alimentary acid is distributed uniformly through its structure, as a result of the acidification being carried out on the dough, and because the acidification treatment after the cooking step ensures a correct acidity or pH to the cooked pasta end product.

[0035] Cooked pasta obtained with the method of the invention may be well preserved for no less than 8 to 12 months at room temperature, or even longer if it is preserved at refrigerator temperature.

[0036] The features and advantages of this invention will be further illustrated by a non-limitative example of the inventive method.

Example

[0037] A dough for alimentary pasta was prepared by mixing together the following ingredients, in percent by weight on the dough weight: 45% semolina of durum fine

flour, 46% water, 9% gluten, and an aqueous solution of glucone-delta-lactone acid having a pH of 2.8 in a preset amount to attain a pH of 3.9 for the dough.

[0038] The kneaded dough had a moisture content of 32%, and was extruded conventionally into conventional spaghetti. The resulting fresh spaghetti were dried conventionally to bring their moisture content down to no more than 11.5%.

[0039] The dried spaghetti were cooked "al dente" for 12 minutes in plenty salted water, and then strained and rinsed with cold water.

[0040] The cooked acidic spaghetti, which had a moisture content of 60%, were dipped in an aqueous solution of glucone- δ -lactone acid having pH 2.2 and a temperature of 22°C for 1-3 minutes.

[0041] At the end of the treatment with acid, the cooked spaghetti had a pH of 4, as measured by conventional analytical methods.

[0042] The cooked acidic spaghetti were taken out of the acid solution, dried, and metered in 200g amounts into suitable trays for food packaging. A preset amount of olive oil was added to each trayful of cooked acidic spaghetti.

[0043] Each trayful of oil-added cooked acidic spaghetti was then sealed in its packaging tray once the atmosphere inside the tray had been modified by removing air and substituting it with nitrogen. The sealed trays containing cooked acidic spaghetti were then pasteurized by microwave treatment.

[0044] The bacterial growth on samples of the spaghetti packaged into the trays was monitored at regular intervals counting from packaging time, through conventional microbiological analysis procedures. The results of these microbiological analyses showed that bacterial growth on the samples tested has been inhibited over the period of 12 months from packaging, and demonstrated a remarkable decrease in average bacterial population. Beyond 12 months after packaging, a slight progressive increase in bacterial population was noticeable which led to eventual decay of the product.

[0045] The above samples of spaghetti were tested for compressive strength on an INSTRON instrument at regular intervals after packaging.

[0046] The results of these tests showed that the compressive strength of the samples of cooked spaghetti was substantially unchanged up to 8 months from packaging, and after 12 months from packaging, the average value was found to be just 5% lower than the value measured before packaging.

Claims

1. A method of producing cooked alimentary pasta, characterized in that it comprises the steps of:

preparing an alimentary dough comprising semolina, water and at least one edible acid;

forming said alimentary dough obtaining fresh acidic pasta having a pH of 3.8 to 4.3 and a moisture content of 30 to 32%;

drying said fresh acidic pasta to a moisture content of 11.0 to 12.5%;

substantially completely cooking said dried pasta obtaining cooked acidic pasta;

treating said cooked acidic pasta with an aqueous solution of an alimentary acid to obtain cooked acidic pasta having a pH not more than 4.1; and

packaging said cooked acidic pasta and subjecting the packaged pasta to thermal sanitation.

2. A method according to Claim 1, wherein said alimentary dough contains gluten in amounts from 4 to 10% by weight on the dough weight, preferably in a 9% amount.
3. A method according to either Claim 1 or 2, wherein said edible acid is provided in the dough in amounts of 0.8 to 1.5% by weight on the dough weight, preferably in a 1% amount.
4. A method according to Claim 3, wherein said edible acid is selected from the group comprising citric acid, fumaric acid, lactic acid, maleic acid, tartaric acid, sulphuric acid, hydrochloric acid, glucone-delta-lactone acid, and phosphoric acids.
5. A method according to Claim 4, wherein said edible acid is glucone-delta-lactone acid.
6. A method according to any of the preceding claims, wherein said cooked acidic alimentary pasta has a moisture content of 53 to 62%.
7. A method according to any of the preceding claims, wherein said step of acid-treating the cooked alimentary pasta is carried out by dipping the cooked acidic alimentary pasta into said aqueous solution of the edible acid for 60 to 180 seconds.
8. A method according to Claim 7, wherein the pH of said aqueous solution of the edible acid is 2.2 to 2.8.
9. A method according to any of the preceding claims, wherein the pH of said cooked acidic pasta is 3.8 to 4.1.
10. A method according to any of the preceding claims, wherein said cooked acidic alimentary pasta is vacuum packaged or packaged under a modified at-

mosphere.

5

10

15

20

25

30

35

40

45

50

55

5



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 42 5101

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	EP 0 968 659 A (NESTLE SA) 5 January 2000 (2000-01-05) * paragraph '0009! * * paragraph '0021! * * paragraph '0036! * * paragraph '0044! *		A23L1/16
A	US 5 332 587 A (FIELD ERIC SYDNEY ET AL) 26 July 1994 (1994-07-26) * column 5, line 13 - line 20 * * column 9, line 18 - line 32 *		
A	PATENT ABSTRACTS OF JAPAN vol. 012, no. 198 (C-502), 8 June 1988 (1988-06-08) & JP 63 000271 A (IZUTSUYA:KK), 5 January 1988 (1988-01-05) * abstract *		
A	EP 0 745 331 A (NESTLE SA) 4 December 1996 (1996-12-04) * example 1 * * page 3, line 7 - page 4, line 29 *		TECHNICAL FIELDS SEARCHED (Int.Cl.7)
A	EP 0 947 143 A (NESTLE SA) 6 October 1999 (1999-10-06) * examples 1,2 * * paragraph '0007! * * paragraph '0008! * * paragraph '0030! *		A23L
D,A	US 4 597 976 A (DOSTER ROBERT C ET AL) 1 July 1986 (1986-07-01) * column 2, line 52 - line 63 * * column 3, line 3 - line 7 * * column 3, line 25 - line 37 *		
-/--			
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 24 July 2002	Examiner Popa, M
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 (03.02) (P04001)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 42 5101

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	US 4 552 772 A (SAITOH HIROSHI ET AL) 12 November 1985 (1985-11-12) * column 1, line 39 - column 2, line 65 * * column 4, line 1 - line 18 * -----		
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 24 July 2002	Examiner Popa, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/82 (P04001)

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 02 42 5101

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24-07-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0968659	A	05-01-2000	EP 0968659 A1	05-01-2000
			AU 3683599 A	13-01-2000
			BR - 9902467 A	16-05-2000
			JP 2000050822 A	22-02-2000
			NZ 335093 A	30-08-1999
			SG 81973 A1	24-07-2001
			US 6187357 B1	13-02-2001
US 5332587	A	26-07-1994	AT 128816 T	15-10-1995
			AT 95989 T	15-11-1993
			AU 639941 B2	12-08-1993
			AU 6293290 A	08-04-1991
			AU 631252 B2	19-11-1992
			AU 6355390 A	08-04-1991
			CA 2040438 A1	02-03-1991
			CA 2065262 A1	02-03-1991
			DE 69004071 D1	25-11-1993
			DE 69004071 T2	03-03-1994
			DE 69023000 D1	16-11-1995
			DE 69023000 T2	21-03-1996
			EP 0415787 A1	06-03-1991
			EP 0415788 A1	06-03-1991
			EP 0489811 A1	17-06-1992
			EP 0441937 A1	21-08-1991
			ES 2079484 T3	16-01-1996
			ES 2047342 T3	16-02-1994
			WO 9103171 A1	21-03-1991
			GR 3018138 T3	29-02-1996
			HU 60113 A2	28-08-1992
			HU 214550 B	28-08-1998
			IE 903186 A1	13-03-1991
			IE 903187 A1	13-03-1991
			JP 5500003 T	14-01-1993
			JP 2654481 B2	17-09-1997
			JP 4501513 T	19-03-1992
			KR 125010 B1	27-11-1997
			NZ 235135 A	28-07-1992
			NZ 235136 A	25-03-1994
			RU 2018235 C1	30-08-1994
			WO 9103169 A1	21-03-1991
			US 5312639 A	17-05-1994
			US 5308632 A	03-05-1994
JP 63000271	A	05-01-1988	NONE	
EP 0745331	A	04-12-1996	US 5599573 A	04-02-1997

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 42 5101

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24-07-2002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0745331 A		AT 207307 T	15-11-2001
		AU 699269 B2	26-11-1998
		AU 5462096 A	12-12-1996
		CA 2177765 A1	01-12-1996
		CN 1140558 A ,B	22-01-1997
		DE 69616150 D1	29-11-2001
		DE 69616150 T2	20-06-2002
		DK 745331 T3	26-11-2001
		EP 0745331 A2	04-12-1996
		ES 2164204 T3	16-02-2002
		JP 8322490 A	10-12-1996
		NO 962218 A	02-12-1996
		NZ 286686 A	26-06-1998
		PT 745331 T	28-03-2002
		SG 73983 A1	18-07-2000
EP 0947143 A	06-10-1999	EP 0953293 A1	03-11-1999
		EP 0947143 A1	06-10-1999
		AU 2139899 A	14-10-1999
		BR 9901207 A	16-05-2000
		CN 1232631 A	27-10-1999
		JP 2000060469 A	29-02-2000
		SG 81958 A1	24-07-2001
		US 6203840 B1	20-03-2001
US 4597976 A	01-07-1986	EP 0239656 A1	07-10-1987
		AU 584413 B2	25-05-1989
		AU 5614186 A	22-10-1987
		CA 1238231 A1	21-06-1988
		GB 2188530 A ,B	07-10-1987
		AT 40937 T	15-03-1989
		DE 3662169 D1	06-04-1989
US 4552772 A	12-11-1985	JP 1019866 B	13-04-1989
		JP 1537354 C	21-12-1989
		JP 57206350 A	17-12-1982
		US 4599238 A	08-07-1986

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82